


# LHC physics : the first 1-2 year(s) ....

Fabiola Gianotti and Michelangelo Mangano  
CERN, PH Department



- ① Physics opportunities at the beginning
- ② Machine start-up scenario
- ③ Which detectors, triggers and performance at the beginning ?  
Construction → test beam → cosmics → first collisions
- ④ How well will we know the physics and the Monte Carlo generators at the beginning ?
- ⑤ Physics goals and potential with the first  $\text{fb}^{-1}$  (a few examples ...)

# ① What can we reasonably expect from the first year(s)?

Some history:

-- Fall 1982: first physics run for UA1 and UA2 at the Sp̄p̄barS

$$L_{\max} = 5 \times 10^{28} \text{cm}^{-2} \text{s}^{-1} \approx 1\% \text{ asymptotic } L$$

$$L_{\text{int}} = 20 \text{nb}^{-1} \text{ in 30 days}$$

outcome: **W/Z discovery, as expected**

ingredients: plenty of kinematical phase-space (ISR was sub-threshold!),  
clear signature, and good hands-on control of backgrounds

-- Summer 1987: first physics run for CDF at the Tevatron

$$L_{\max} = 5 \times 10^{28} \text{cm}^{-2} \text{s}^{-1} \approx 1\% \text{ nominal } L$$

$$L_{\text{int}} = 20 \text{nb}^{-1} \text{ in 30 days}$$


outcome: **nothing exciting, as expected**

why: not enough phase-space, given the strong constraints on new physics  
already set by UA1/UA2!

In the region of the UA1 limit the production cross-section at the Tevatron was only a factor of 10-20 larger

By the time of CDF startup, the SppS had already logged enough luminosity to rule out a possible observation at the Tevatron within the first  $100\text{nb}^{-1}$


It took 2 more years (and  $4\text{pb}^{-1}$ ) for CDF to improve ( $m_{\text{top}} > 77\text{ GeV}$ ) the UA1 limits (in spite of the fact that by '89, and with  $5\text{pb}^{-1}$ , had only improved to  $60\text{ GeV}$  - UA2 eventually went up to  $69\text{ GeV}$ ). This is the consequence of much higher bg's at the Tevatron, and of the steep learning curve for such a complex analysis



At the start of LHC, the situation will resemble much more that at the beginning of UA1/UA2:

The phase-space for the Tevatron will have totally saturated the search boundary for most phenomena, at a level well below the LHC initial reach: seen from the LHC, the Tevatron will look like the ISR as seen from the SppS!

Rates  $10^3$  times larger in the region of asymptotic Tevatron reach




1% of  $L_{\text{max}}$  for the LHC,  
(as in SppS and Tevatron  
early runs),  
close to  $L_{\text{max}}$  for Tevatron

(assume a 1% signal efficiency)

N.B.: rates for gluino  
production are roughly a  
factor of 10 larger than  
for HQs



Similar considerations hold for jets, where few days of data will probe quarks at scales beyond the overall Tevatron CM energy!



Fine, we have phase-space, we have rates. But should we truly expect something to show up at scales reachable early on?

LEP's heritage is a strong confirmation of the SM, and at the same time an apparent paradox:

on one side  $m(H)=117+45-68$ ; on the other, SM radiative corrections give

$$\delta m_H^2 = \frac{6G_F}{\sqrt{2}\pi^2} (m_t^2 - \frac{1}{2}m_W^2 - \frac{1}{4}m_Z^2 - \frac{1}{4}m_H^2)\Lambda^2 \sim (115\text{GeV})^2 \left(\frac{\Lambda}{400\text{GeV}}\right)^2$$

How can counterterms artificially conspire to ensure a cancellation of their contribution to the Higgs mass?

The existence of new phenomena at a scale not much larger than 400 GeV appears necessary to enforce such a cancellation in a natural way!

The accuracy of the EW precision tests at LEP, on the other hand, sets the scale for "generic new physics" (parameterized in terms of dim-5 and dim-6 effective operators) at the level of few-to-several TeV.

This sets very strong constraints on the nature of this possible new physics: to leave unaffected the SM EW predictions, and at the same time to play a major role in the Higgs sector.

**Supersymmetry offers one such possible solution**

In Supersymmetry the radiative corrections to the Higgs mass are not quadratic in the cutoff, but logarithmic in the size of SUSY breaking (in this case  $M_{\text{stop}}/M_{\text{top}}$ ):


$$m_h^2 < m_Z^2 + \frac{3G_F}{\sqrt{2}\pi^2} m_t^4 \left[ \ln \left( \frac{M_S^2}{m_t^2} \right) + x_t^2 \left( 1 - \frac{x_t^2}{12} \right) \right] \quad \text{with} \quad M_S^2 \equiv \frac{1}{2}(M_{t_1}^2 + M_{t_2}^2) \quad X_t \equiv A_t - \mu \cot \beta$$

$$x_t \equiv X_t/M_S$$

For  $M_{\text{susy}} < 2\text{TeV}$

$m_h^{\text{max}} \simeq 122 \text{ GeV}$ , if top-squark mixing is minimal,

$m_h^{\text{max}} \simeq 135 \text{ GeV}$ , if top-squark mixing is maximal



The current limits on  $m_H$  point to  $M(\text{lightest stop}) > 600 \text{ GeV}$ . Pushing the SUSY scale towards the TeV, however, forces fine tuning in the EW sector, reducing the appeal of SUSY as a solution to the Higgs mass naturalness:

$$\delta m_Z^2 \sim (90 \text{ GeV})^2 \left( \frac{M_S}{230 \text{ GeV}} \right)^2 \ln \frac{\Lambda_{UV}}{M_S}$$

In other words, the large value of  $m_H$  shows that room is getting very tight now for SUSY, at least in its "minimal" manifestations. **This makes the case for an early observation of SUSY at the LHC quite compelling, and worth investing into!**

For some people the room left is too tight. Some skepticism on SUSY has emerged, and a huge effort of looking for alternatives has began few years back, leading to a plethora of new ideas (Higgsless-models, Little Higgs, extra-dimensions, etc)

Some of these ideas lead to rather artificial structures, where the problem of the Higgs naturalness is shifted to slightly higher scales, via the introduction of a new sector of particles around the TeV.

The observation of new phenomena within the first few yrs of run, in these cases, is not guaranteed (nor is it asymptotically)

Few of these scenarios offer the appeal of Supersymmetry, with its clear predictions (calculability), and connections with the other outstanding problems of the Standard Model (Dark Matter, Flavour, CP violation)

# Dark matter constraints on neutralinos: a CMSSM example

old:  
 $0.1 < \Omega_\chi h^2 < 0.3$

new WMAP



$0.094 < \Omega_\chi h^2 < 0.129$

$$\Omega_\chi h^2 \sim m_\chi n_\chi \Rightarrow$$



upper limit on  $\Omega_\chi$  requires:

+ small  $m_\chi$ , or


+ fast/efficient annihilation, a strong constraint on spectrum (to allow, e.g.,  $\chi\chi \rightarrow h$  at threshold or  $\chi\tau \rightarrow \gamma\tau$ )  $\sim$



J. Ellis et al, hep-ph/0303043




In the CMSSM the measurement of  $m_{1/2}$  and  $m_0$  (resp.  $m_\chi$  and  $m_{\text{slep}}$ ) will fix almost uniquely  $\tan\beta$



**Proving the direct and unambiguous link between cosmology, DM and SUSY would be, perhaps even more than the Higgs discovery, the flagship achievement of the LHC**

The search for Supersymmetry is in my view the single most important task facing the LHC experiments in the early days. In several of its manifestations, SUSY provides very clean final states, with large rates and potentially small bg's.



Given the big difficulty and the low rates characteristic of Higgs searches in the critical domain  $m_H < 135 \text{ GeV}$ , I feel that the **detector and physics commissioning should be optimized towards the needs of SUSY searches rather than light-Higgs** (I implicitly assume that for  $m_H > 140$  Higgs searches will be almost straightforward and will require proper understanding of only a limited fraction of the detector components -- e.g. muons)

The early determination of the scale at which new physics manifests itself will have important consequences for the planning of facilities beyond the LHC (LC? CLIC? nufact? Flavour factories? Underground Dark Matter searches?).

The LHC will have no competition in the search for new physics, so in principle there is no rush. But the future of the field will greatly benefit from a quick feedback on SUSY and the rest!



## ② Machine start-up scenario → see L.Rossi

(from Chamonix XII Workshop, January 2003)


~ 400 dipoles delivered  
~ 300 cold-tested



- ~ April 2007 : start machine cool-down followed by machine commissioning (mainly with single beam)
- ~ Summer 2007 : two beams in the machine → first collisions
  - 43 + 43 bunches,  $L=6 \times 10^{31} \text{ cm}^{-2} \text{ s}^{-1}$  (possible scenario; tuning machine parameters)
  - pilot run: 936+936 bunches (75 ns → no electron cloud),  $L > 5 \times 10^{32}$
  - 2-3 month shut-down ?
  - 2808 + 2808 bunches (bunch spacing 25 ns),  $L$  up to  $\sim 2 \times 10^{33}$  (goal of first year)  
→ ~ 7 months of physics run

A lot of uncertainties in this plan (QRL !) → here show potential vs integrated luminosity from  $\sim 100 \text{ pb}^{-1} / \text{expt}$  to  $\sim 10 \text{ fb}^{-1} / \text{expt}$

### ③ Which detectors the first year(s)?




RPC over  $|\eta| < 1.6$  (instead of  $|\eta| < 2.1$ )  
4<sup>th</sup> layer of end-cap chambers missing

Pixels and end-cap ECAL  
installed during first shut-down

2 pixel layers/disks instead of 3

TRT acceptance over  $|\eta| < 2$   
(instead of  $|\eta| < 2.4$ )




Both experiments:  
deferrals of high-level Trigger/DAQ processors  
→ LVL1 output rate limited to  
    ~ 50 kHz CMS (instead of 100 kHz)  
    ~ 35 kHz ATLAS (instead of 75 kHz)

Impact on physics visible but acceptable

Main loss : B-physics programme strongly reduced (single  $\mu$  threshold  $p_T \rightarrow 14-20$  GeV)

# Which detector performance at day one ?

A few examples and educated guesses based on test-beam results and simulation studies





	Expected performance day 1	Physics samples to improve (examples)
ECAL uniformity e/ $\gamma$ scale	$\sim 1\%$ (ATLAS), 4% (CMS) 1-2 % ?	Minimum-bias, $Z \rightarrow ee$ $Z \rightarrow ee$
HCAL uniformity Jet scale	2-3 % < 10%	Single pions, QCD jets $Z (\rightarrow ll) + 1j, W \rightarrow jj$ in $t\bar{t}$ events
Tracking alignment	20-500 $\mu\text{m}$ in $R\phi$ ?	Generic tracks, isolated $\mu$ , $Z \rightarrow \mu\mu$

Ultimate statistical precision achievable after few days of operation. Then face systematics ....  
E.g. : tracker alignment : 100  $\mu\text{m}$  (1 month)  $\rightarrow$  20 $\mu\text{m}$  (4 months)  $\rightarrow$  5  $\mu\text{m}$  (1 year) ?


# Steps to achieve the detector goal performance

- Stringent construction requirements and quality controls (piece by piece ...)
- Equipped with redundant calibration/alignment hardware systems
- Prototypes and part of final modules extensively tested with test beams (allows also validation of Geant4 simulation)
- In situ calibration at the collider (accounts for material, global detector, B-field, long-range mis-calibrations and mis-alignments) includes :
  - cosmic runs : end 2006-beg 2007 during machine cool-down
  - beam-gas events, beam-halo muons during single-beam period
  - calibration with physics samples (e.g.  $Z \rightarrow ll$ ,  $t\bar{t}$ , etc.)







## Example of this procedure : ATLAS electromagnetic calorimeter



Pb-liquid argon sampling calorimeter  
with Accordion shape, covering  $|\eta| < 2.5$




$H \rightarrow \gamma\gamma$  : to observe signal peak on top of huge  $\gamma\gamma$  background need mass resolution of  $\sim 1\%$   $\rightarrow$  response uniformity (i.e. total constant term of energy resolution)  $\leq 0.7\%$  over  $|\eta| < 2.5$




① Construction phase (e.g. mechanical tolerances):

287 GeV electron response variation with Pb thickness from '93 test-beam data




1% more lead in a cell → 0.7% response drop  
 → to keep response uniform to 0.2-0.3%, thickness of Pb plates must be uniform to 0.5% (~ 10 μm)

Thickness of all 1536 absorber plates (1.5m long, 0.5m wide) for end-cap calorimeter measured with ultrasounds during construction




② **Beam tests** of 4 (out of 32) barrel modules and 3 (out of 16) end-cap modules:

1 barrel module:  
 $\Delta\eta \times \Delta\phi = 1.4 \times 0.4$   
 $\equiv \sim 3000$  channels




Scan of a barrel module with 245 GeV  $e^-$



Uniformity over "units" of size  
 $\Delta\eta \times \Delta\phi = 0.2 \times 0.4 : \sim 0.5\%$   
 400 such units over the full ECAL



③ Check calibration with **cosmic muons**:




From full simulation of ATLAS (including cavern, overburden, surface buildings) + measurements with scintillators in the cavern:




Through-going muons (hits in ID + top and bottom muon chambers)	~ 25 Hz
Pass by origin ( $ z  < 60$ cm, $R < 20$ cm, hits in ID)	~ 0.5 Hz
Useful for ECAL calibration ( $ z  < 30$ cm, $E_{\text{cell}} > 100$ MeV, $\sim 90^\circ$ )	~ 0.5 Hz

→ ~  $10^6$  events in ~ 3 months of data taking  
→ enough for initial detector shake-down  
(catalog problems, gain operation experience,  
some alignment/calibration, detector synchronization, ...)






Precision of ECAL readout calibration system : 0.25%.  
 But :  $\eta$ -dependent differences between calibration and physics signals



→ can be checked with cosmic muons



From studies with test-beam muons:  
 can check (and correct) calorimeter response variation vs  $\eta$  to 0.5% in < 3 months of cosmic runs

Note : not at level of ultimate calibration uniformity ( $\sim 0.25\%$ ) but already a good starting point

④ First collisions : calibration with  $Z \rightarrow ee$  events

rate  $\sim 1$  Hz at  $10^{33}$ ,  $\sim$  no background, allows ECAL standalone calibration

$$c_{\text{tot}} = c_L \oplus c_{\text{LR}}$$



$c_L \approx 0.5\%$  demonstrated at the test-beam over units  $\Delta\eta \times \Delta\phi = 0.2 \times 0.4$   
 $c_{\text{LR}} \equiv$  long-range response non-uniformities from unit to unit (400 total)  
 (module-to-module variations, different upstream material, etc.)

Use  $Z \rightarrow ee$  events and  $Z$ -mass constraint to correct long-range non-uniformities.

From full simulation :  $\sim 250 e^\pm$  / unit needed to achieve  $c_{\text{LR}} \leq 0.4\% \rightarrow c_{\text{tot}} = 0.5\% \oplus 0.4\% \leq 0.7\%$

$\sim 10^5 Z \rightarrow ee$  events (few days of data taking at  $10^{33}$ )

Nevertheless, let's consider the worst (unrealistic ?) scenario : no corrections applied

- $c_L = 1.3\%$
- $c_{\text{LR}} = 1.5\%$

measured "on-line" non-uniformity of individual modules  
 no calibration with  $Z \rightarrow ee$

}  $\longrightarrow$   $c_{\text{tot}} \approx 2\%$



conservative : implies very poor knowledge of upstream material (to factor  $\sim 2$ )

$H \rightarrow \gamma\gamma$  significance  $m_H \sim 115$  GeV degraded by  $\sim 25\%$   
 $\rightarrow$  need 50% more L for discovery

## ④ How well will we know LHC physics on day one (before data taking starts) ?

- \* DY processes
- \* top X-sections
- \* bottom X-sections
- \* jet X-sections
- \* Higgs X-sections

# W/Z cross-sections



- Test of QCD to NNLO: potential accuracy  $\sim 2\%$  on  $\sigma_{\text{tot}}$
  - Luminosity monitor
  - Probe of PDF's
- => In view of incomplete detector coverage, need to ensure that the potential NNLO accuracy is reflected in the calculation of acceptancies. The realization of a QCD NNLO event generator, however, will still take few years. Is it required?

Cuts A  $\rightarrow |\eta^{(e)}| < 2.5, p_T^{(e)} > 20 \text{ GeV}, p_T^{(\nu)} > 20 \text{ GeV}$



Cuts B  $\rightarrow |\eta^{(e)}| < 2.5, p_T^{(e)} > 40 \text{ GeV}, p_T^{(\nu)} > 20 \text{ GeV}$

	LO	LO+HW	NLO	MC@NLO
<b>Cuts A</b>	0.5249	-7.7% 0.4843	0.4771	+1.5% 0.4845
	↓5.4%		↓7.0%	↓6.3%
<b>Cuts A, no spin</b>	0.5535		0.5104	0.5151
<b>Cuts B</b>	0.0585	+208% 0.1218	0.1292	+2.9% 0.1329
	↓29%		↓16%	↓18%
<b>Cuts B, no spin</b>	0.0752		0.1504	0.1570

Theory OK to 2% + 2%(PDF)

Similar accuracy for high-mass DY (bg, as well as signal, for massive Z'/W')

# tt cross-section



$$\sigma_{tt}^{\text{FNAL}} = 6.5 \text{ pb} (1 \pm 5\%_{\text{scale}} \pm 7\%_{\text{PDF}})$$

Scale unc:  $\pm 12\%_{\text{NLO}} \Rightarrow \pm 5\%_{\text{NLO+NLL}}$

$$\sigma_{tt}^{\text{LHC}} = 840 \text{ pb} (1 \pm 5\%_{\text{scale}} \pm 3\%_{\text{PDF}})$$

$\Delta\sigma = \pm 6\% \Rightarrow \Delta m = \pm 2 \text{ GeV}$

# Recent overview of ATLAS strategy and results for $m_{top}$ : hep-ph/0403021

## Channels considered:

- + (W $\rightarrow$ l $\nu$ )+4 jets, 2 b tags
- + high-pT top, t $\rightarrow$ 3 jets
- + (W $\rightarrow$ l $\nu$ ) (W $\rightarrow$ l $\nu$ ) + bb
- +  $m_{(l\text{-}psi)}$  in events with B $\rightarrow$ psiX

Source of error in GeV	Lepton+jets inclusive sample	Lepton+jets large clusters sample	Dilepton	All jets high pT sample
Energy scale				
Light jet energy scale	0.2	-	-	0.8
b-jet energy scale	0.7	-	0.6	0.7
Mass scale calibration	-	0.9	-	-
UE estimate	-	1.3	-	-
Physics				
Background	0.1	0.1	0.2	0.4
b-quark fragmentation	0.1	0.3	0.7	0.3
Initial state radiation	0.1	0.1	0.1	0.4
Final state radiation	-	0.1	0.6	2.8
PDF	-	-	1.2	-

## Need a strategy for validation of the MC input models:


+ UE modeling and subtraction

+ validation of FSR effects:

\* jet fragmentation properties, jet energy profiles

\* how do we validate emission off the top quark in the high-pt top sample?

\* b fragmentation function






# bb cross-sections

OK, but theoretical systematics still large:

+/-35% at low pt  
 +/-20% for  $p_T \gg m_b$

In view of the recent run II results from CDF, more validation required.

To verify the better predictivity at large pt, need to perform measurements in the region 30-80 GeV, and above (also useful to study properties of high-Et b jets, useful for other physics studies)




# Higgs cross-sections

NNLO available for dominant  $gg \rightarrow H$  process  
=> almost as accurate as DY

PDF uncert sufficient for day-1 business, but improvements necessary for high-lum x-sec studies (=> to measure couplings)

(Djouadi & Ferrag, hep-ph/0310209)







# Jet cross-sections


Theoretical syst uncertainty  
at NLO  $\sim \pm 20\%$

PDF uncert (mostly  $g(x)$ ) growing at large  $x$




# DO, run I data

## Cone jets (R=0.7)



## $k_T$ jets (D=1)



**Puzzling discrepancy at low  $E_T$ , in view of the fact that at NLO rates for cone-jets with  $R=0.7$  and  $k_T$  jets with  $D=1$  are equal to within 1%**

**OK at high- $E_T$**

# Main sources of syst uncertainties (CDF, run I)

At high  $E_T$  the syst is dominated by the response to high  $p_T$  hadrons (beyond the test beam  $p_T$  range) and fragmentation uncertainties

Out to which  $E_T$  will the systematics allow precise cross-section measurements at the LHC?

Out to which  $E_T$  can we probe the jet structure (multiplicity, fragm function)?

NB: stat for Z+jet or gamma+jet runs out before  $E_T \sim 500$  GeV




Table 8: Rates for  $L_{int} = 10 fb^{-1}$  for different intervals of  $P_t^Z$  and  $\eta^Z$  ( $P_{tCUT}^{clust} = 10 GeV/c$ ,  $P_{tCUT}^{out} = 10 GeV/c$  and  $\Delta\phi \leq 15^\circ$ ).

$P_t^Z$ (GeV/c)	$\Delta\eta^Z$   intervals						all   $\eta^Z$
	0.0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-5.0	0.0-5.0
40 – 50	4594	5425	6673	7267	6732	4796	35486
50 – 60	3128	3509	4297	4570	3976	2000	21471
60 – 70	2253	2443	2855	2934	2229	851	13567
70 – 80	1580	1734	1948	1786	1307	341	8692
80 – 90	1152	1148	1267	1236	824	170	5790
90 – 100	741	859	812	808	523	59	3802
100 – 110	582	590	594	546	305	36	2657
110 – 120	384	428	451	412	226	8	1905
120 – 140	523	582	562	531	293	12	2503
140 – 170	392	380	368	341	190	4	1675
170 – 200	170	186	162	170	63	2	756
200 – 240	111	103	99	91	40	0	444
240 – 300	71	51	44	48	20	0	238

Z+jet


$P_{tCUT}^{clust} = 5 GeV/c$  and  $\Delta\phi \leq 15^\circ$ .

(GeV/c)									all $\eta^\gamma$
	0.0-0.4	0.4-0.7	0.7-1.1	1.1-1.5	1.5-1.9	1.9-2.2	2.2-2.6	0.0-2.6	
40 – 50	102656	107148	100668	103903	103499	116674	126546	761027	
50 – 60	43905	41729	41074	45085	42974	47640	50310	312697	
60 – 70	18153	18326	19190	20435	20816	19432	23650	140005	
70 – 80	9848	10211	9963	10166	9951	11397	10447	71984	
80 – 90	5287	5921	5104	5823	5385	6067	5923	39509	
90 – 100	2899	3033	3033	3326	3119	3265	3558	22234	
100 – 120	2908	3091	2995	3305	3133	3282	3429	22143	
120 – 140	1336	1359	1189	1346	1326	1499	1471	9525	
140 – 160	624	643	626	674	706	614	668	4555	
160 – 200	561	469	557	555	519	555	557	3774	
200 – 240	187	176	186	192	187	185	151	1264	
240 – 300	103	98	98	98	100	92	74	665	
300 – 360	34	34	33	32	31	27	20	212	
40 – 360	188517	192274	184734	194957	191761	210742	226819	1389484	


gamma+jet

# The structure of the underlying event

Mounting experimental evidence (R.Field, CDF) that the UE is the result of **multiple semi-hard (minijet-like) interactions**



## “Associated” PTsum Density PYTHIA Tune A vs HERWIG



- Extrapolation from Tevatron to LHC is hard, as it relies on the understanding of the unitarization of the minijet cross-section
- The mini-jet nature of the UE implies that the particle and energy flows are not uniformly distributed within a given event: can one do better than the standard uniform, constant, UE energy subtraction?
- Studies of MB and UE should be done early on, at very low luminosity, to remove the effect of overlapping pp events:
  - MB triggers
  - low- $E_T$  jet triggers

## 5 Physics goals and potential in the first year (a few examples ...)

Channels (examples ...)	Events to tape for $10 \text{ fb}^{-1}$ (per experiment)
$W \rightarrow \mu \nu$	$7 \times 10^7$
$Z \rightarrow \mu \mu$	$1.1 \times 10^7$
$t\bar{t} \rightarrow W b W b \rightarrow \mu \nu + X$	$0.08 \times 10^7$
QCD jets $p_T > 150$	$\sim 10^7$
Minimum bias	$\sim 10^7$
$\tilde{g}\tilde{g}$ $m = 1 \text{ TeV}$	$10^3 - 10^4$

~ few PB of data per year per experiment → challenging for software and computing (esp. at the beginning ...)

} assuming 1% of trigger bandwidth



Already in first year, large statistics expected from:

- known SM processes → understand detector and physics at  $\sqrt{s} = 14 \text{ TeV}$
- several New Physics scenarios

Note: overall event statistics limited by  $\sim 100 \text{ Hz}$  rate-to-storage

$\sim 10^7$  events to tape every 3 days assuming 30% data taking efficiency



## Goal # 1

Understand and calibrate detector and trigger in situ using well-known physics samples

- e.g. -  $Z \rightarrow ee, \mu\mu$  tracker, ECAL, Muon chambers calibration and alignment, etc.  
-  $t\bar{t} \rightarrow b\bar{t} bjj$   $10^3$  evts/day after cuts  $\rightarrow$  jet scale from  $W \rightarrow jj$ , b-tag perf., etc.

Understand basic SM physics at  $\sqrt{s} = 14$  TeV  $\rightarrow$  first checks of Monte Carlos

(hopefully well understood at Tevatron and HERA)

- e.g. - measure cross-sections for e.g. minimum bias, W, Z,  $t\bar{t}$ , QCD jets (to  $\sim 10-20\%$ ),  
look at basic event features, first constraints of PDFs, etc.  
- measure top mass (to 5-7 GeV)  $\rightarrow$  give feedback on detector performance

Note : statistical error negligible after few weeks run

## Goal # 2

Prepare the road to discovery:

- measure backgrounds to New Physics : e.g.  $t\bar{t}$  and W/Z+ jets (omnipresent ...)
- look at specific "control samples" for the individual channels:  
e.g.  $t\bar{t}jj$  with  $j \neq b$  "calibrates"  $t\bar{t}bb$  irreducible background to  $t\bar{t}H \rightarrow t\bar{t}bb$


## Goal # 3

Look for New Physics potentially accessible in first year (e.g. Z', SUSY, some Higgs ? ...)



## Example of initial measurement : top signal and top mass



- Use gold-plated  $t\bar{t} \rightarrow bW bW \rightarrow bl\nu bjj$  channel
- Very simple selection:
  - isolated lepton ( $e, \mu$ )  $p_T > 20$  GeV
  - exactly 4 jets  $p_T > 40$  GeV
  - no kinematic fit
  - no b-tagging required (pessimistic, assumes trackers not yet understood)
- Plot invariant mass of 3 jets with highest  $p_T$



Time	Events at 10 <sup>33</sup>	Stat. error $\delta M_{\text{top}}$ (GeV)	Stat. error $\delta\sigma/\sigma$
1 year	3x10 <sup>5</sup>	0.1	0.2%
1 month	7x10 <sup>4</sup>	0.2	0.4%
1 week	2x10 <sup>3</sup>	0.4	2.5%

- top signal visible in few days also with simple selections and no b-tagging
- cross-section to ~ 20% (10% from luminosity)
- top mass to ~7 GeV (assuming b-jet scale to 10%)
- get feedback on detector performance :
  - $m_{\text{top}}$  wrong  $\rightarrow$  jet scale ?
  - gold-plated sample to commission b-tagging


Fit signal and background (top width fixed to 12 GeV) → extract cross-section and mass



Can we see a  $W \rightarrow jj$  peak?

Select the 2 jets with highest  $p_T$   
(better ideas well possible ...)




$W$  peak visible in signal, no peak in background



# Introduce b-tagging ...


ATLAS 150 pb<sup>-1</sup>

Bkgd composition changes: combinatorial from top itself becomes more and more important




## What about early discoveries ?

An easy case : a new resonance decaying into  $e^+e^-$ , e.g. a  $Z'$   $\rightarrow ee$  of mass 1-2 TeV



An intermediate case : SUSY



A difficult case : a light Higgs ( $m \sim 115$  GeV)




An "easy case" :  $Z'$  of mass 1-2 TeV with SM-like couplings

$Z' \rightarrow ee, \text{SSM}$

Mass	Expected events for $10 \text{ fb}^{-1}$ (after all cuts)	$\int \mathcal{L} dt$ needed for discovery (corresponds to 10 observed evts)
1 TeV	$\sim 1600$	$\sim 70 \text{ pb}^{-1}$
1.5 TeV	$\sim 300$	$\sim 300 \text{ pb}^{-1}$
2 TeV	$\sim 70$	$\sim 1.5 \text{ fb}^{-1}$


- signal rate with  $\int \mathcal{L} dt \sim 0.1\text{-}1 \text{ fb}^{-1}$  large enough up to  $m \approx 2 \text{ TeV}$  if "reasonable"  $Z'ee$  couplings
- dominant Drell-Yan background small  
( $< 15$  events in the region  $1400\text{-}1600 \text{ GeV}$ ,  $10 \text{ fb}^{-1}$ )
- signal as mass peak on top of background

$Z \rightarrow ll + \text{jet}$  samples and DY needed for E-calibration and determination of lepton efficiency



# An intermediate case : SUPERSYMMETRY

Large  $\tilde{q}\tilde{q}, \tilde{q}\tilde{g}, \tilde{g}\tilde{g}$  cross-section  $\rightarrow \approx 100$  events/day at  $10^{33}$  for  $m(\tilde{q}, \tilde{g}) \sim 1$  TeV  
 Spectacular signatures  $\rightarrow$  SUSY could be found quickly




Using multijet +  $E_T^{miss}$  (most powerful and model-independent signature if R-parity conserved)

~ one year at  $10^{34}$ :  
up to ~2.5 TeV



~ one year at  $10^{33}$ :  
up to ~2 TeV

~ one month at  $10^{33}$ :  
up to ~1.5 TeV



Measurement of sparticle masses likely requires > 1 year. However ...

Peak position correlated to  $M_{\text{SUSY}} \equiv \min(m(\tilde{q}), m(\tilde{g}))$



ATLAS

From  $M_{\text{eff}}$  peak  $\rightarrow$  first/fast measurement of SUSY mass scale to  $\approx 20\%$  (10 fb<sup>-1</sup>, mSUGRA)

Detector/performance requirements:

- quality of  $E_T^{\text{miss}}$  measurement (calorimeter inter-calibration/linearity, cracks)
  - $\rightarrow$  apply hard cuts against fake MET and use control samples (e.g.  $Z \rightarrow ll + \text{jets}$ )
- "low" Jet /  $E_T^{\text{miss}}$  trigger thresholds for low masses at overlap with Tevatron region ( $\sim 400 \text{ GeV}$ )

Backgrounds will be estimated using data (control samples) and Monte Carlo:


Background process (examples ...)	Control samples (examples ...)
$Z (\rightarrow \nu\nu) + \text{jets}$ $W (\rightarrow \tau\nu) + \text{jets}$ $t\bar{t} \rightarrow b\bar{t}b\bar{t}j$ QCD multijets	$Z (\rightarrow ee, \mu\mu) + \text{jets}$ $W (\rightarrow e\nu, \mu\nu) + \text{jets}$ $t\bar{t} \rightarrow b\bar{t}b\bar{t}$ lower $E_T$ sample

Can estimate background levels also varying selection cuts (e.g. ask 0,1,2,3 leptons ...)

A lot of data will most likely be needed!

normalization point

normalise MC to data at low  $E_T^{\text{miss}}$  and use it to predict background at high  $E_T^{\text{miss}}$  in "signal" region





Hard cuts against fake  $E_T^{\text{miss}}$  :

- reject beam-gas, beam-halo, cosmic
- primary vertex in central region
- reject event with  $E_T^{\text{miss}}$  vector along a jet or opposite to a jet
- reject events with jets in cracks
- etc. etc.



# Can we trust the current estimates of bg rates?





$N_{\text{jet}} \geq 4$

$E_{T(1,2)} > 100 \text{ GeV}$

$E_{T(3,4)} > 50 \text{ GeV}$


$\text{MET} > \max(100, M_{\text{eff}}/4)$

$M_{\text{eff}} = \text{MET} + \sum E_{Tj}$

“Correct” bg shape  
indistinguishable  
from signal shape!

Bg breakdown:

- QCD jets
- $t\bar{t}$
- ▼  $Z \rightarrow \nu\nu$
- ▲  $W \rightarrow l\nu$



Indeed the  $Z \rightarrow \nu\nu$  bg appears to be underestimated by a factor 10–50! It will dominate the highMET tail, and could be measured in  $Z \rightarrow ee + \text{jets}$

Use  $Z \rightarrow ee$  + multijets, apply same cuts as MET analysis but replace MET with  $ET(e^+e^-)$


Extract  $Z \rightarrow \nu\nu$  bg using, bin-by-bin:  
 $(Z \rightarrow \nu\nu) = (Z \rightarrow ee) B(Z \rightarrow \nu\nu)/B(Z \rightarrow ee)$

Assume that the SUSY signal is of the same size as the bg, and evaluate the luminosity required to determine the  $Z \rightarrow \nu\nu$  bg with an accuracy such that:

$$N_{\text{susy}} > 3 \text{ sigma}$$

where


$$\text{sigma} = \sqrt{N(Z \rightarrow ee)} * B(Z \rightarrow \nu\nu)/B(Z \rightarrow ee)$$



=> several hundred  $\text{pb}^{-1}$  are required. They are sufficient if we believe in the MC shape (and only need to fix the overall normalization). Much more is needed if we want to keep the search completely MC independent


**How to validate the estimate of the MET from resolution tails in multijet events??**

# A difficult case: a light Higgs $m_H \sim 115 \text{ GeV}$



$m_H \sim 115 \text{ GeV}$      $10 \text{ fb}^{-1}$

total  $S/\sqrt{B} \approx 4^{+2.2}_{-1.3}$



ATLAS	$H \rightarrow \gamma\gamma$	$ttH \rightarrow ttbb$	$qqH \rightarrow qq\tau\tau$ ( $ll + l\text{-had}$ )
S	130	15	$\sim 10$
B	4300	45	$\sim 10$
$S/\sqrt{B}$	2.0	2.2	$\sim 2.7$




$\uparrow$  K-factors  $\equiv \sigma(\text{NLO})/\sigma(\text{LO}) \approx 2$  not included

Full GEANT simulation, simple cut-based analyses

## Remarks:

Each channel contributes  $\sim 2\sigma$  to total significance  $\rightarrow$  **observation of all channels important to extract convincing signal in first year(s)**

The 3 channels are complementary  $\rightarrow$  robustness:



- different production and decay modes
- different backgrounds
- different detector/performance requirements:
  - **ECAL crucial for  $H \rightarrow \gamma\gamma$**  (in particular response uniformity) :  $\sigma/m \sim 1\%$  needed
  - **b-tagging crucial for  $ttH$**  : 4 b-tagged jets needed to reduce combinatorics
  - **efficient jet reconstruction over  $|\eta| < 5$  crucial for  $qqH \rightarrow qq\tau\tau$**  : forward jet tag and central jet veto needed against background



Note : -- **all require "low" trigger thresholds**

E.g.  $ttH$  analysis cuts :  $p_T(l) > 20 \text{ GeV}$ ,  $p_T(\text{jets}) > 15\text{-}30 \text{ GeV}$

-- **all require very good understanding (1-10%) of backgrounds**


If  $m_H > 180 \text{ GeV}$  : early discovery may be easier with  $H \rightarrow 4l$  channel

Luminosity needed for  $5\sigma$  discovery (ATLAS+CMS)



- $H \rightarrow WW \rightarrow l\nu l\nu$  : high rate ( $\sim 100 \text{ evts/expt}$ ) but no mass peak  $\rightarrow$  not ideal for early discovery ...
  - $H \rightarrow 4l$  : low-rate but very clean : narrow mass peak, small background
- Requires: --  $\sim 90\%$   $e, \mu$  efficiency at low  $p_T$  (analysis cuts :  $p_T^{1,2,3,4} > 20, 20, 7, 7, \text{ GeV}$ )  
 --  $\sigma / m \sim 1\%$ , tails  $< 10\%$   $\rightarrow$  good quality of  $E, p$  measurements in ECAL and tracker

## A crucial role in these measurements is played by the vector boson fusion process:




To suppress the bg's, typical analyses require, in addition to the decay products of the H, the following:

- \* Two jets with large  $M(jj)$ , one forward and one backward (typically  $|\eta| > 2.5$ )
- \* A veto on central jets ( $|\eta| < 2.5$ ), justified by the lack of colour exchange between the two hadrons, leading to a rapidity gap





Standard analyses of jet veto efficiency use ME calculations for  $qq \rightarrow Hqq$ , with the central jet generated via a parton shower. Angular ordering in the parton shower prevents emission of central jets, and a bad underestimate of the signal events with a central jet!

Exact Hqq+jet  
Naive Hqq+shower



Central jets in Hqq events are therefore usually assumed to originate from additional multiple collisions. This is quite true at high luminosity, but not at  $10^{33}$



Correct determination of veto efficiency for signal is not just important to establish the best threshold for discovery, but to evaluate the signal cross-section after discovery!

**No data from the Tevatron or elsewhere allow today to validate our estimates of central-jet emission in VBF processes. This needs to be done, possibly using the low-luminosity data where fake jets due to multiple interactions are strongly reduced.**

(table from F.Cerutti)

Channel	Main background	S/B	background systematics for $5\sigma$	Proposed technique/comments
H $\rightarrow\gamma\gamma$	Irreduc. $\gamma\gamma$ Reducible $\gamma j$	2-3%	0.4%	Side-bands stat Err $\sim 0.5\%$ for 30-100 fb $^{-1}$
$t\bar{t}H$ H $\rightarrow bb$	$t\bar{t}jj$	30%	6%	Mass side-bands Anti b-tagged $t\bar{t}jj$ ev. Under study
H $\rightarrow ZZ^*\rightarrow 4$ lep	ZZ $\rightarrow 4l$ and $\tau\tau ll$	300-600%	60%	Mass side-bands Stat Err $< 30\%$ 30fb $^{-1}$
H $\rightarrow WW^*\rightarrow ll\nu\nu$	WW*, $tW$	30-50%	6%	No mass peak Bkg enriched region ? Study to be performed
VBF channels In general	Rejection QCD/EW	Study forward jet tag and central jet veto		Use EW ZZ and WW leptonic Study to be performed
VFB H $\rightarrow WW$	$t\bar{t}$ , WW, Wt	50-200%	10%	Study Z,W,WW and $t\bar{t}$ plus jets
VBF H $\rightarrow\tau\tau$	Zjj, $t\bar{t}$	50-400%	10%	Missing Et calibration Z $\rightarrow\tau\tau$ (mass tails ?) Study to be performed
MSSM (bb)H/A $\rightarrow\tau\tau$	Z $\rightarrow\tau\tau$ , Wj	25% $t\bar{g}\beta=15$ $M_A=300$	5%	Mass side-bands Stat Err $\sim 5\%$ 30fb $^{-1}$
MSSM (bb)H/A $\rightarrow \mu\mu$	Z/ $\gamma^*\rightarrow\mu\mu$	12% $t\bar{g}\beta=15$ $M_A=150$	$\sim 2\%$	Mass side-bands Stat Err $\sim 2\%$ 30fb $^{-1}$

# Conclusions



- LHC has potential for major discoveries already in the first year (months ?) of operation  
Event statistics : 1 day at LHC at  $10^{33}$   $\equiv$  1 year at previous machines for SM processes  
SUSY may be discovered "quickly", light Higgs more difficult ... and what about surprises ?
- Machine luminosity performance will be the crucial issue in first 1-2 years
- Experiments: lot of emphasis on test beams and on construction quality checks  
→ results indicate that detectors "as built" should give good starting-point performance.
- However: lot of data (and time ...) will be needed at the beginning to:
  - commission the detector and trigger in situ (and the software !!! ...)
  - reach the performance needed to optimize the physics potential
  - understand standard physics at  $\sqrt{s} = 14$  TeV and compare to MC predictions  
[ Tevatron (and HERA) data crucial to speed up this phase ... ]
  - measure backgrounds to possible New Physics (with redundancy from several samples ...)
- Efficient/robust commissioning with physics data in the various phases (cosmics, one-beam period, first collisions, ...), as well as solid preparation of MC tools, are our next challenges.  
Both are crucial to reach quickly the "discovery-mode" and extract a convincing "early" signal

# Back-up slides

- Variation of the jet energy scale to infer systematics
  - Bjet scale: 0.92 – 0.96 – 1.00 – 1.04 – 1.08
  - Light scale: 0.94 – 0.98 – 1.00 – 1.02 – 1.04
- Determine  $M_{\text{top}}$  and  $\sigma(\text{top})$ 
  - ‘Raw’, i.e. no correction for jet scale
  - ‘Corrected’, i.e. apply percentage difference of W-peak to the reconstructed top
  - Not granted  $M_{jj}$  gives correct MW, i.e. for hard FSR events...
- Dependence on top mass reduced by scaling with W:
  - Rms of top masses:
    - Raw: 6.2 GeV
    - Scaled: 1.2 GeV
  - Note: Here simple rescaling of Top mass – not of the jet-energies themselves!
- Large dependence  $\sigma(\text{top})$  on jet energy scale
  - Via event selection.

### Scale variations:

5 scales for each of the three generators (MC@NLO Pythia Herwig) and for MC@NLO with 2 times background added




# Commissioning ID with cosmics and beam gas (preliminary ideas ...)

## Cosmics : O (1Hz) tracks in Pixels+SCT+TRT


- useful statistics for debugging readout, maps of dead modules, etc.
- check relative position Pixels/SCT/TRT and of ID wrt ECAL and Muon Spectrometer
- first alignment studies: may achieve statistical precision of  $\sim 10 \mu\text{m}$  in parts of Pixels/SCT
- first calibration of R-t relation in straws

## Beam-gas :

- $\sim 25 \text{ Hz}$  of reconstructed tracks with  $p_T > 1 \text{ GeV}$  and  $|z| < 20 \text{ cm}$   
→  $> 10^7$  tracks (similar to LHC events) in 2 months
- enough statistics for alignment in "relaxed" environment → exceed initial survey precision of  $10\text{-}100 \mu\text{m}$



standard ATLAS patt. rec.  
(no optimisation for cosmics ...)





LVL1 menus and rates (indicative only ...)

**ATLAS**


**CMS**

$L = 2 \cdot 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$	Threshold (GeV)	Rate (kHz)	Threshold (GeV)	Rate (kHz)
Inclusive muon	20	0.8	14	2.7
Two muons	6	0.2	3	0.9
Inclusive electron	25	12.0	29	3.3
Two electrons	15	4.0	17	1.3
1 Jet, 3 Jet, 4 Jet	200, 90, 65	0.6	177, 86, 70	3.0
Jet + $E_T^{\text{miss}}$	60-60	0.4	88-46	2.3
tau + $E_T^{\text{miss}}$	25-30	2.0		
Inclusive tau			86	2.2
Two taus			59-59	1.0
Electron + Jet			21-45	0.8
Others (pre-scaled, calibration, ...)		5.0		0.9
<b>Total</b>		<b>~ 25</b> (no safety margin)		<b>~16</b> (factor ~3 safety margin)

→ B-physics programme strongly reduced (e.g.  $B \rightarrow J/\psi (\rightarrow ee) K_S^0$ , hadronic channels)

# Which trigger ?

CMS,  $L = 2 \times 10^{33}$



LVL1	Channel	Threshold [GeV] $\epsilon = 95\%$	Rate [kHz]
	Inclusive isolated e/ $\gamma$	29	3.3
	Di-electrons/di-photons	17	1.3
	Inclusive isolated muon	14	2.7
	Di-muons	3	0.9
	Single-tau / two-taus	86/59	2.2/1.0
	1-jet, 3-jets, 4-jets	177 , 86 , 70	3.0
	Jet * $E_{T,miss}$	88 * 46	2.3
	Min-bias (Calibration)		0.9
	<b>Total</b>		<b>16 kHz</b>

## HLT (to tape)

Channel	Threshold [GeV] $\epsilon = 90...95\%$	Rate [Hz]
1 e, 2 e	29 , 17 + 17	34
1 $\gamma$ , 2 $\gamma$	80 , 40 + 25	9
1 $\mu$ , 2 $\mu$	19 , 7 + 7	29
1 $\tau$ , 2 $\tau$	86, 59 + 59	4
1jet OR 3jet OR 4	657 , 247, 113	9
Jet * $E_{T,miss}$	180 + 123	5
Calibration, Other		~17
<b>Total (purity ~50%)</b>		<b>~105 Hz</b>

~ 50 kHz with x3 safety

- LVL1 rate limited by staging of HLT processors
- HLT rate by cost of offline computing (1 PB/year)
- Should preserve guiding principles of LHC trigger !  
 Inclusive approach to the "unknown", safe overlap with Tevatron reach, avoid biases from exclusive selections, margin for offline optimization and QCD uncertainties, enough bandwidth for calibration/control triggers (esp. at beginning !)

- HLT/DAQ deferrals limit available networking and computing for HLT → limit LVL1 output rate
- Large uncertainties on LVL1 affordable rate vs money (component cost, software performance, etc.)

Selections (examples ...)	LVL1 rate (kHz) L= 1 × 10 <sup>33</sup> no deferrals	LVL1 rate (kHz) L= 2 × 10 <sup>33</sup> no deferrals	LVL1 rate (kHz) L= 2 × 10 <sup>33</sup> with deferrals An example for illustration...
MU6,8,20	23	→ 19	→ 0.8
2MU6	---	0.2	0.2
EM20i,25,25	11	→ 12	→ 12
2EM15i,15,15	2	4	4
J180,200,200	0.2	0.2	0.2
3J75,90,90	0.2	0.2	0.2
4J55,65,65	0.2	0.2	0.2
J50+xE50,60,60	0.4	0.4	0.4
TAU20,25,25 +xE30	2	2	2
MU10+EM15i	---	0.1	0.1
Others (pre-scaled, etc.)	5	5	5
Total	~ 44	~ 43	~ 25

LVL1 designed for 75 kHz  
→ room for factor ~ 2 safety

Likely max affordable rate,  
no room for safety factor

### ③ Which data samples ?

Total trigger rate to storage at  $2 \times 10^{33}$   
reduced from  $\sim 540$  Hz (HLT/DAQ TP, 2000)  
to  $\sim 200$  Hz (now)

High-Level-Trigger output



Selection (examples ...)	Rate to storage at $2 \times 10^{33}$ (Hz)	Physics motivations (examples ...)
$e25i, 2e15i$	$\sim 40$ (55% W/b/c $\rightarrow eX$ )	Low-mass Higgs ( $\tau\tau H, H \rightarrow 4\ell, qq\tau\tau$ )
$\mu20i, 2\mu10$	$\sim 40$ (85% W/b/c $\rightarrow \mu X$ )	W, Z, top, New Physics ?
$\gamma60i, 2\gamma20i$	$\sim 40$ (57% prompt $\gamma$ )	$H \rightarrow \gamma\gamma$ , New Physics (e.g. $X \rightarrow \gamma\gamma$ $m_X \sim 500$ GeV) ?
$j400, 3j165, 4j110$	$\sim 25$	Overlap with Tevatron for new $X \rightarrow jj$ in danger ...
$j70 + xE70$	$\sim 20$	SUSY : $\sim 400$ GeV squarks/gluinos
$\tau35 + xE45$	$\sim 5$	MSSM Higgs, New Physics (3 <sup>rd</sup> family !) ? More difficult high L
$2\mu6 (+ m_B)$	$\sim 10$	Rare decays $B \rightarrow \mu\mu X$
Others	$\sim 20$	Only 10% of total !
(pre-scaled, exclusive, ...)		
<b>Total</b>	<b><math>\sim 200</math></b>	<b>No safety factor included.</b>

Best use of spare capacity when  $L < 2 \times 10^{33}$  being investigated

"Signal" (W,  $\gamma$ , etc.) :  $\sim 100$  Hz

# Impact also on high- $p_T$ physics : ~ no safety margin left

Main impact expected on light Higgs

To include factor ~ 2 safety (e.g. QCD cross-sections likely higher than expected) should limit rate to ~ 10 kHz (!):

- must raise EM trigger thresholds, e.g. :
  - from 2EM15i (4 kHz) to 2EM20i (1 kHz) → what about light H → 4e ( $p_T > 20, 20, 7, 7$  GeV) ?
  - from EM25i (12 kHz) to EM30i (4.5 kHz)
- and/or must use less inclusive selections
  - what about total rate when summing all possible channels ? E.g.
  - what about biases (e.g. final states with low- $p_T$  jets, small  $E_T^{miss}$ ) ?
  - what about unknown discovery physics ?
- must decrease pre-scaled/control triggers (note : should rather be increased if higher thresholds and more exclusive menus)

$ttH \rightarrow l\nu bb + X$   $m_H = 120$  GeV

Thresholds (GeV)	Normalised S/ $\sqrt{B}$
$p_T(e) > 20, p_T(\mu) > 20$	1
$p_T(e) > 25, p_T(\mu) > 20$	0.98
$p_T(e) > 30, p_T(\mu) > 20$	0.96
$p_T(e) > 30, p_T(\mu) > 30$	0.92
$p_T(e) > 35, p_T(\mu) > 25$	0.92

Physics TDR (reference)

with deferrals depending on e.g. QCD cross-sections


EM25i + 2J30	4 kHz	} OR= 9 kHz
EM25i + xE15	7 kHz	
EM25i + tau35	?	
EM25i + ?	?	
Total	> 9 kHz	← qqH → $\tau\tau$

not much smaller than EM25i (12 kHz) !

Note : ~ 8% loss from pixel staging not included

# Jet triggers already at the limit for overlap with Tevatron

E.g. : New particles decaying into two jets



CDF/D0 reach for 15 fb<sup>-1</sup>:

$m \sim 700-1200 \text{ GeV}$  (95% C.L.)

→ Jacobian peak at  $p_T(\text{jet}) \sim 350-600 \text{ GeV}$

ATLAS :

single-jet trigger threshold :  $p_T = 400 \text{ GeV}$



di-jet trigger threshold :  $p_T = 350 \text{ GeV} ?$



Relevant issues for early discovery:


- J70+xE70 thresholds for unprescaled triggers
- enough pre-scaled lower-threshold triggers to normalize B
- quality of  $E_T^{miss}$  measurement (calorimeter inter-calibration, cracks)

Cracks can be monitored with  $Z (\rightarrow ll) + jets$   
 B rejection tools:  
 $E_T^{miss}$  isolation, removal of jets in cracks



“Poor” initial calorimeter calibration may increase trigger rates → impact on low-mass SUSY  
 Uncorrected non-compensation simulated by + 20% enhancement of EM scale → + 50% rate for  $E_T^{miss} > 80$  GeV

## What about dead channels ?



Requirement : fraction of dead channels  $< 0.3\%$   
Measurements of the final assembled ECAL  
(at warm and cold) gave :  $\sim 0.1\%$  of dead channels

## Summary of physics impact of staging initial detector

Staged items	Main impact during first run on	Effect
1 pixel layer	$t\bar{t}H \rightarrow t\bar{t}bb$	~8% loss in significance
Gap scintillator	$H \rightarrow 4e$	~8% loss in significance
MDT	$A/H \rightarrow 2\mu$	~5% loss in significance for $m \sim 300 \text{ GeV}$
Trigger processors	B-physics → High- $p_T$ physics →	program jeopardised no safety margin (e.g. for EM triggers)

Requires 10-15% more integrated luminosity to compensate.

Complete detector needed at high luminosity:

- robust pattern recognition (efficiency, fakes rate) in the presence of pile-up and radiation background
- muon measurement
- powerful b-tag
- robustness against detector aging and  $L > 10^{34}$
- precise measurements (e.g. light Higgs) may require low trigger thresholds

} at (very) high  $p_T$

# Data samples for calibration and control


- ❶ Well-known, clean processes from standard trigger menu: e.g.  $t\bar{t}$ ,  $Z \rightarrow \ell\bar{\ell}$
- ❷ Additional lower-thresholds samples needed (esp. at the beginning) → pre-scaled triggers

- **Minimum-bias events:** pp interaction properties, MC tuning, LVL1 efficiency, radiation background in Muon chambers, etc.
- **QCD jets ( $20 \leq E_T \leq 400 \text{ GeV}$ ):** QCD cross-sections and MC tuning, trigger efficiency, calorimeter inter-calibration, jet algorithms, background to Higgs, SUSY, etc.
- **Inclusive  $e^\pm$   $p_T > 10 \text{ GeV}$ :** trigger efficiency, ECAL calibration, ID alignment,  $E/p$ ,  $e^\pm$  reconstruction at low- $p_T$ , etc.
- **efficiency,  $\mu^\pm$  reconstruction at low- $p_T$ , calorimeters, ID alignment, etc.**

These are only few examples ...

$\sim 10^7$  events per sample

Rate :  
 $\sim 10 \text{ Hz/sample}$  first weeks  
 $\sim \text{few Hz/sample}$  under normal operation



$E_T^{\text{miss}}$  resolution vs  $\Sigma E_T$  using minimum-bias and QCD jets (full GEANT3 simulation)

$\geq 10\%$  of total rate

# Which physics the first year(s) ?

Expected event rates at production in ATLAS or CMS at  $L = 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$

Process	Events/s	Events for $10 \text{ fb}^{-1}$	<u>Total statistics collected</u> at previous machines by 2007
$W \rightarrow e\nu$	15	$10^8$	$10^4$ LEP / $10^7$ Tevatron
$Z \rightarrow ee$	1.5	$10^7$	$10^6$ LEP
$t\bar{t}$	1	$10^7$	$10^4$ Tevatron
$b\bar{b}$	$10^6$	$10^{12} - 10^{13}$	$10^9$ Belle/BaBar ?
H $m=130 \text{ GeV}$	0.02	$10^5$	?
$\tilde{g}\tilde{g}$ $m=1 \text{ TeV}$	0.001	$10^4$	---
Black holes $m > 3 \text{ TeV}$ ( $M_D=3 \text{ TeV}, n=4$ )	0.0001	$10^3$	---



Already in first year, large statistics expected from:

- known SM processes → understand detector and physics at  $\sqrt{s} = 14 \text{ TeV}$
- several New Physics scenarios

Systematic error on  $m_{top}$  (TDR performance,  $10 \text{ fb}^{-1}$ )


<i>Source of uncertainty</i>	<i>Hadronic part</i> $\delta M_{Top} \text{ (GeV)}$	<i>Kinematic fit</i> $\delta M_{Top} \text{ (GeV)}$	<i>Comments</i>  1% error 1% error ( $\epsilon_b = -0.006$ ) - ( $\epsilon_b = -0.035$ ) 20%(ON-OFF) 20%(ON-OFF)
Light jet energy scale	0.9	0.2	
b-jet energy scale	0.7	0.7	
b-quark frag.	0.1	0.1	
ISR	0.1	0.1	
FSR	1.9	0.5	
Combinatorial Bkg	0.4	0.1	
<b>Total</b>	<b>2.3</b>	<b>0.9</b>	


Initial performance : uncertainty on b-jet scale expected to dominate

b-jet scale uncertainty	$\delta m \text{ (top)}$
1%	0.7 GeV
5%	3.5 GeV
10%	7 GeV

Cfr: 10% on q-jet scale +  $m_W$  (PDG)  $\rightarrow$  3 GeV on  $m(\text{top})$

Initial  $\delta m \text{ (top)} \sim 5-7 \text{ GeV} ?$





$m_H \sim 130 \text{ GeV} \quad 10 \text{ fb}^{-1}$

$l = e, \mu$

	$H \rightarrow \gamma\gamma$	$qqH \rightarrow qq\tau\tau$ ( $ll + l\text{-had}$ )	$H \rightarrow 4l$	$qqH \rightarrow qqWW$
S	120	$\sim 8$	$\sim 5$	18
B	3400	$\sim 6$	$< 1$	15
S/√B	2.0	$\sim 2.7$	2.8	3.9

↑ K-factor  $\approx 2$  not included ↑

total  $S/\sqrt{B} \approx 6$

- 4 complementary channels for physics and for detector requirements
- $S/\sqrt{B} < 3$  per channel (except  $qqWW$  counting channel) → observation of all channels important in first year

•  $H \rightarrow 4l$  low rate but very clean: small background, narrow mass peak

Detector requirements:

- $\geq 90\%$   $e, \mu$  efficiency at low  $p_T$  (analysis cuts :  $p_T^{1,2,3,4} > 20, 20, 7, 7, \text{ GeV}$ )
- in particular low di-lepton LVL1 thresholds




Channel	Main background	S/B	background systematics for 5 $\sigma$	Proposed technique/comments
H $\rightarrow\gamma\gamma$	Irreduc. $\gamma\gamma$ Reducible $\gamma j$	2-3%	0.4%	Side-bands stat Err $\sim 0.5\%$ for 30-100 fb $^{-1}$
$t\bar{t}H$ H $\rightarrow bb$	$t\bar{t}jj$	30%	6%	Mass side-bands Anti b-tagged $t\bar{t}jj$ ev. Under study J.Cammin
H $\rightarrow ZZ^* \rightarrow 4$ lep	ZZ $\rightarrow 4l$ and $\tau\tau ll$	3-6	60%	Mass side-bands Stat Err $< 30\%$ 30fb $^{-1}$
H $\rightarrow WW^* \rightarrow ll\nu\nu$	WW*, $tW$	30-50%	6%	No mass peak Bkg enriched region ? Study to be performed
VBF channels In general	Rejection QCD/EW	Study forward jet tag and central jet veto		Use EW ZZ and WW leptonic Study to be performed
VFB H $\rightarrow WW$	$t\bar{t}$ , WW, Wt	50-200%	10%	Bkg. enriched samples with discr. Variables Study to be performed
VBF H $\rightarrow \tau\tau$	Zjj, $t\bar{t}$	50-400%	10%	Missing Et calibration Z $\rightarrow \tau\tau$ (mass tails ?) Study to be performed
MSSM (bb)H/A $\rightarrow \tau\tau$	Z $\rightarrow \tau\tau$ , Wj	25% $t\bar{g}\beta=15$ MA=300	5%	Mass side-bands Stat Err $\sim 5\%$ 30fb $^{-1}$
MSSM (bb)H/A $\rightarrow \mu\mu$	Z/ $\gamma^* \rightarrow \mu\mu$	12% $t\bar{g}\beta=15$ MA=150	$\sim 2\%$	Mass side-bands Stat Err $\sim 2\%$ 30fb $^{-1}$

# MSSM Higgs bosons : $h, H, A, H^\pm$


$h$  : similar to SM Higgs over most of the allowed region

$m_h < 135 \text{ GeV}$   
 $m_A \approx m_H \approx m_{H^\pm}$  at large  $m_A$



--  $bbA, bbH, H^\pm$  cross-section  $\sim tg^2\beta$   
 -- best sensitivity from  $A/H \rightarrow \tau\tau, H^\pm \rightarrow \tau\nu$   
 (not easy the first year ...)  
 --  $A/H \rightarrow \mu\mu$  experimentally easier  
 (esp. at the beginning)

Requires non-ultimate b-tagging (one jet),  
 and non-ultimate tracking resolution ( $A/H$   
 intrinsic width non negligible)




# MSSM Higgs bosons $h, H, A, H^\pm$

$$m_h < 135 \text{ GeV}$$

$$m_A \approx m_H \approx m_{H^\pm} \text{ at large } m_A$$

5 $\sigma$  discovery curves

September 2002




--  $A, H, H^\pm$  cross-section  $\sim \text{tg}^2\beta$

-- best sensitivity from  $A/H \rightarrow \tau\tau, H^\pm \rightarrow \tau\nu$   
(not easy the first year ...)


--  $A/H \rightarrow \mu\mu$  experimentally easier  
(esp. at the beginning)

Measurement of  $\text{tg} \beta$



- Large variety of channels and signatures accessible
- $bbA/H \rightarrow 4b$  is more difficult than at the Tevatron (because of huge QCD background)

September 2002




Here  $\geq 5\sigma$   
discovery of  
 $bbA/H \rightarrow 4b$   
possible at  
Tevatron with  
 $15 \text{ fb}^{-1}$

# SUSY mass scale ( $\sim$ model-independent)


D.Tovey

$M_{\text{SUSY}}$  (GeV)




$M_{\text{SUSY}}$

$M_{\text{eff}}$  (GeV)




$M_{\text{eff}}$  (GeV)

% precision on  $M_{\text{SUSY}}$  vs  $M_{\text{SUSY}}$



\* 10 fb<sup>-1</sup>  
 ○ 100 fb<sup>-1</sup>  
 ● 300 fb<sup>-1</sup>



conservative!

Intrinsic spread from model parameters  
 (infinite statistics, no experimental error):

$\sim 2\%$  mSUGRA


$\sim 10\%$  constrained MSSM

Including experimental uncertainties ( $\sim 50\%$  from background subtraction,  $\sim 1.5\%$  from E-scale):

$\leq 20\%$  ( $10\%$ ) mSUGRA for 10 (100) fb<sup>-1</sup>

$\leq 60\%$  ( $30\%$ ) constrained MSSM for 10 (100) fb<sup>-1</sup>

## Precision on measured SUSY cross-section vs $M_{\text{SUSY}}^{\text{eff}}$





Including experimental uncertainties :

- ≤ 30% mSUGRA for 300 fb<sup>-1</sup>
- ≤ 80% constrained MSSM for 300 fb<sup>-1</sup>



## Theoretical SUSY cross-section vs $M_{\text{SUSY}}^{\text{eff}}$



Z'


Quick discovery, assuming SM couplings (SSM)

mass	$\sigma \times \text{BR}(Z \rightarrow e e)$ in peak	events, $10 \text{ fb}^{-1}$
1 TeV	360 fb	3600
1.5 TeV	64 fb	640
2.0 TeV	15.7 fb	157

present limits:  
690 GeV (direct),  
1500 GeV(EW fit)

Allows to compare and test different detector components for high energy particles:  $ee, \mu\mu, \tau\tau, bb, jj$




Z--> ll + jets samples needed for E calibration





# Expected rates of beam-gas events

Vertex z-position	Rate (Hz)	Total (2 months, $\epsilon=30\%$ )
$\pm 23$ m	$1.2 \cdot 10^5$	$2.1 \cdot 10^{11}$
$\pm 3$ m	$1.6 \cdot 10^4$	$2.4 \cdot 10^{10}$
$\pm 20$ cm	$1.1 \cdot 10^3$	$1.6 \cdot 10^9$
$\pi^\pm$ $p_T > 1$ GeV inside $\pm 3$ m	$1.0 \cdot 10^3$	$1.5 \cdot 10^9$
$\gamma$ $p_T > 1$ GeV inside $\pm 3$ m	$0.3 \cdot 10^3$	$5.6 \cdot 10^8$



## Expected rates of beam-halo muons

- Rates for initial period scaled from high-luminosity rates by assuming  $3 \times 10^{10}$  p per bunch and 43 bunches  $\rightarrow \sim 200$  times lower current
- Expected optics and vacuum for commissioning period not included yet (need input from machine people)  $\rightarrow$  these results are very preliminary
- Total rates are for two months of single-beam with 30% data taking efficiency
- Simple definition of "useful tracks" : 2-3 segments in MDT, 3-4 disks in ID end-cap

Detector	Rate (B-field off)	Total (B-field off)	Rate (B-field on)	Total (B-field on)
MDT barrel	15 Hz	$2.5 \cdot 10^7$	72 Hz	$1.5 \cdot 10^8$
MDT end-cap	145 Hz	$2.5 \cdot 10^8$	135 Hz	$2.5 \cdot 10^8$
Pixel/SCT	1.8/17 Hz	$3 \cdot 10^6 / 3 \cdot 10^7$	2/19 Hz	$3 \cdot 10^6 / 3 \cdot 10^7$
EM $E > 5 \text{ GeV}$	2 Hz	$3.5 \cdot 10^6$	1 Hz	$1.7 \cdot 10^6$
Tile/HEC $E > 20 \text{ GeV}$	1.7/1.2 Hz	$2.9/2.1 \cdot 10^6$	1.6/0.9 Hz	$2.8/1.6 \cdot 10^6$

Very preliminary